arXiv:1402.3408v1 [math.DS] 14 Feb 2014

ON THE QUOTIENT SPACE OF A GENERALIZED ACTION OF A GENERALIZED GROUP

HASAN MALEKI, MOHAMMADREZA MOLAEI

Department of Mathematics, Shahid Bahonar University of Kerman, Kerman,

Iran

e-mails: maleki_hasan@yahoo.com; mrmolaei@uk.ac.ir

Abstract

In this paper by using of generalized groups and their generalized actions, we define and study the notion of T-spaces. Moreover, we study properties of the quotient space of a T-space and we present the conditions that imply to the Hausdorff property for it. Moreover, we study the maps between two T-spaces and we consider the notion of T-transitivity.

Key words: Generalized action; Transitivity; T-space; Quotient space

1 Introduction

Groups appears in many branches of mathematics such as number theory, geometry and theory of lie groups and also they can find plenty applications even in physics and chemistry. Generalized groups are an interesting and fascinating extension of groups. They have been introduced in 1999 [9]. In a group, there is a unique identity element but in a generalized group, each element has a spacial identity element. This kind of structure appears in genetic codes [12]. The role of identity as a mapping made many new

important challenges. Generalized groups have been applied in genetic [1, 12], geometry [11] and dynamical systems [13]. The notion of generalized action [8] is an extension of the notion of group actions [3, 6]. This notion has been studied first in 1999 [8, 10, 12]. The generalized action has been applied by the other researchers [4, 5].
Let us recall the definition of a generalized group or completely simple semi group [2]. A *generalized group* is a semigroup T with the following conditions:

(i) For each $t \in T$ there exists a unique $e(t) \in T$ such that $t \cdot e(t) = e(t) \cdot t = t$;

(ii) For each $t \in T$ there exists $s \in T$ such that $s \cdot t = t \cdot s = e(t)$.

One can easily prove that: each t in T, has a unique inverse in T. The inverse of t is denoted by t^{-1}. Moreover; for given $t \in T$, $e(t) = e(t^{-1})$ and $e(e(t)) = e(t)$. Let T and S be tow generalized groups. A map $f : T \to S$ is called a *homomorphism* if $f(st) = f(s) \cdot f(t)$ for every $s, t \in T$. We refer to [12] for more details. The geometrical viewpoint of completely simple semigroups are taken in [14]. Here we study more properties of the generalized action of a generalized group on a topological spaces. In the next section we present the notion of T-spaces. We consider the quotient space of a T-space and we deduce the conditions that imply to the Hausdorff property for it. In the last section the maps between two T-spaces and T-transitivity are considered.

2 T-Spaces

Let us to begin this section by recalling the definition of topological generalized group.

Definition 2.1. [1, 10] A *topological generalized group* is a Hausdorff topological space T which is endowed with a generalized group structure such that the generalized group operations $m_1 : T \to T$ defined by $t \mapsto t^{-1}$ and $m_2 : T \times T \to T$ by $(s, t) \mapsto s \cdot t$ are continuous maps and

$$e(s \cdot t) = e(s) \cdot e(t). \tag{1}$$

As Shown in [12], every topological generalized group T is a disjoint union of topological groups. Moreover, the mapping $e : T \to T$ defined by $t \mapsto e(t)$, is a continuous map. Note that in a topological generalized group, the property (1) implies that

$$e(s) \cdot e(t) \cdot e(s) = e(s), \ \forall t, s \in T. \tag{2}$$

This property will be used frequently.

Theorem 2.2. *If T is a topological generalized group,then*

$$(s \cdot t)^{-1} = e(s) \cdot t^{-1} \cdot s^{-1} \cdot e(t)$$

Proof.

$$\begin{aligned}
(s \cdot t)(e(s) \cdot t^{-1} \cdot s^{-1} \cdot e(t)) &= s \cdot (t \cdot e(s) \cdot t^{-1} \cdot s^{-1} \cdot e(t)) \\
&= s \cdot (t \cdot e(t) \cdot e(s) \cdot e(t) \cdot t^{-1} \cdot s^{-1} \cdot e(t)) \\
&= s \cdot (t \cdot e(t) \cdot t^{-1} \cdot s^{-1} \cdot e(t)) \\
&= s \cdot (e(t) \cdot s^{-1} \cdot e(t)) \\
&= (s \cdot e(t) \cdot s^{-1}) \cdot e(t) \\
&= (s \cdot e(s) \cdot e(t) \cdot e(s) \cdot s^{-1}) \cdot e(t) \\
&= (s \cdot e(s) \cdot s^{-1}) \cdot e(t) \\
&= e(s) \cdot e(t) = e(s \cdot t).
\end{aligned}$$

We also have $(e(s) \cdot t^{-1} \cdot s^{-1} \cdot e(t)) \cdot (s \cdot t) = e(s \cdot t)$. So $(s \cdot t)^{-1} = e(s) \cdot t^{-1} \cdot s^{-1} \cdot e(t)$. □

Example 2.3. If T is the topological space

$$\mathbb{R}^2 - \{(0,0)\} = \{re^{i\theta} | \quad r > 0 \quad \text{and} \quad 0 \leqslant \theta < 2\pi\}$$

with the Euclidean metric, then T with the multiplication

$$(r_1 e^{i\theta_1}) \cdot (r_2 e^{i\theta_2}) = r_1 r_2 e^{i\theta_2} \tag{3}$$

is a topological generalized group. We have $e(re^{i\theta}) = e^{i\theta}$ and $(re^{i\theta})^{-1} = \frac{1}{r} e^{i\theta}$. So we can see the identity set $e(T)$ is the unit circle S^1. However, T is not a topological group.

Definition 2.4. Suppose that X is a topological space and T is a topological generalized group. A *generalized action* of T on X is a continuous map $\lambda : T \times X \longrightarrow X$ with the following properties:

(i) $\lambda(s, \lambda(t, x)) = \lambda(s \cdot t, x)$, for $s, t \in T$ and $x \in X$;

(ii) If $x \in X$, then is $e(t) \in T$ such that $\lambda(e(t), x) = x$.

Henceforth, $\lambda(t, x)$ will be denoted by tx. For the more details on the generalized action, one can see [5, 8]. Now let us define the notion of T-spaces.

Definition 2.5. A *T-space* is a triple (X, T, λ) where X is a Hausdorff topological space, T is a topological generalized group and $\lambda : T \times X \longrightarrow X$ is a generalized action of T on X.

Definition 2.6. If (X, T, λ) is a T-space and $x \in X$, then:

(i) $T_x = \{t \in T | \quad tx = x\}$ is called the *stabilizer* of x in T;

(ii) $T(x) = \{tx | \quad t \in T\}$ is called *T-orbit* of x in X.

For $x \in X$, T_x is a generalized subgroup of T [5]. For $x \in X$, if the stabilizer T_x is trivial (i.e. $T_x = \{e(t)\}$ for some $t \in T$), we say that x is a *regular* point, otherwise, it is called a *singular* point. The *singular* set of X, denoted by $\sum_X$, is the set of singular points of X. We define two maps $\theta_t : X \to X$ and $\rho_x : T \to X$, by $\theta_t(x) = tx$ and $\rho_x(t) = tx$, respectively, where $t \in T$ and $x \in X$. θ_t and ρ_x are continuous maps. Clearly, $T(x) = \rho_x(T)$ and $T_x = (\rho_x)^{-1}(x)$. So T_x is a closed subset of T and then we can say it is a closed generalized subgroup of T. Furthermore, T_x is a topological generalized subgroup of T [5].

By the action λ of a T-space (X, T, λ), we can define the following equivalence relation on X:

$$x \sim y \text{ if and only if there is } t \in T \text{ such that } tx = y.$$

Now, we consider the quotient space $X/_\sim$ and by the projection map $\pi : X \to X/_\sim$, we define a topology on $X/_\sim$ such that π is a continuous map. In fact $U \subseteq X/_\sim$ is open if $\pi^{-1}(U)$ is open in X.

Henceforth, we will use of the notation $X/_T$ for the topological quotient space $X/_\sim$.

Definition 2.7. Suppose (X, T, λ) is a T-space. Then the generalized action λ is called:

(i) *effective* if for any two distinct $s, t \in T$, there exists $x \in X$ such that $sx \neq tx$;

(ii) *transitive* if for given $x, y \in X$, there exists $t \in T$ such that $tx = y$;

(iii) *free* if for each $x \in X$, there exists precisely one $t \in T$ such that $tx = x$;

(iv) *regular* if it is transitive and free.

λ is transitive if and only if $T(x) = X$, for each $x \in X$ and λ is free if and only if $T_x = \{e(t)\}$, for each $x \in X$.

Example 2.8. Every topological generalized group T acts on itself by the multiplication of T : $st = s \cdot t$ for all $s, t \in T$. Note that this action need not be free, for instance, the action $xy = y$ as the multiplication of T implies $T_x = T$, for $x \in X$.

Example 2.9. Every topological generalized group T acts on itself by the multiplication $st = e(s) \cdot t$.

Example 2.10. Let T be $\mathbb{R} - \{0\}$ with the Euclidean metric. T with the multiplication $x \cdot y = x$ is a topological generalized group that if $x \in T$, then $e(x) = x^{-1} = x$. T acts on itself with this multiplication. This action is regular.

We recall that if T is a generalized group, X is a set and $S = \{\varphi^t \mid \varphi^t : X \to X$ is a mapping and $t \in T\}$, then the triple (X, S, T) is called a *complete semidynamical system* if:

(i) $\varphi^{t_1} \circ \varphi^{t_2} = \varphi^{t_1 \cdot t_2}$, for all $t_1, t_2 \in T$;

(ii) For given $x \in X$, there is $\varphi^t \in D$ such that x is a fixed point of φ^t.

We see that each T-space (X, T, λ) generates a complete semidynamical system (X, S, T) where

$$S = \{\theta_t : X \to X \mid \theta_t(x) = tx, \text{ for } x \in X \text{ and } t \in T\}.$$

As shown in [5], if $e(T) \subseteq T_x$, for each $x \in X$, then S with the multiplication $\theta_s \circ \theta_t = \theta_{st}$ is a topological generalized group. In this case, for each $\theta_t \in S, e(\theta_t) = \theta_{e(t)}$ and $(\theta_t)^{-1} = \theta_{t^{-1}}$.

Theorem 2.11. *If $e(T) \subseteq T_x$, for each $x \in X$, then each θ_t is a homeomorphism.*

Proof. Every θ_t is a continuous map on X. Now we claim that each θ_t is one to one, onto and has the inverse $\theta_{t^{-1}}$. If $\theta_t(x) = \theta_t(y)$, then $tx = ty$, and so $t^{-1}tx = e(t)x = x = y = e(t)y = t^{-1}ty$. Hence θ_t is one to one. On the other hand, for $x \in X$, there exists $t^{-1}x \in X$ such that $\theta_t(t^{-1}x) = tt^{-1}x = e(t)x = x$. Thus θ_t is also onto. If $t \in T$ and $x \in X$, then

$$\theta_t \circ \theta_{t^{-1}}(x) = \theta_{tt^{-1}}(x) = \theta_{e(t)}(x) = e(t)x = x.$$

The last equity follows from the fact $e(T) \subseteq T_x$. In the same way

$$\theta_{t_{-1}} \circ \theta_t(x) = x.$$

Therefore, every θ_t is a homeomorphism. □

Theorem 2.12. *Let (X, T, λ) be a T-space. If T is compact and $e(T) \subseteq T_x$ for each $x \in X$, then $\lambda : T \times X \to X$ is a closed map.*

Proof. Assume C is a closed subset of $T \times X$ and $x \in X$ is a limit point of $\lambda(C)$. So there exists a sequence $\{(t_i, x_i)\}$ in C such that $\lambda(t_i, x_i) = t_i x_i$ converges to x. As T is compact, then there is a subsequence of $\{t_i\}$ such that converges to a t in T. We rename that subsequence be $\{t_i\}$. T is a topological generalized group, so the map $m_1 : T \to T$, defined by $t \mapsto t^{-1}$ is continuous. This implies that $\{t_i^{-1}\}$ converges to t^{-1}. λ is also continuous, $\lambda(t_i^{-1}, t_i x_i)$ converges to $\lambda(t^{-1}, x)$, so $\{e(t_i)x_i\}$ converges to $t^{-1}x$. But for each $x \in X$, $e(T) \subseteq T_x$, thus $e(t_i)x_i = x_i$ and consequently, x_i converges to $t^{-1}x$. So the sequence $\{(t_i, x_i)\}$ in C converges to $(t, t^{-1}x)$. Since C is a closed subset of $T \times X$, then $(t, t^{-1}x) \in C$. Therefore, $\lambda(t, t^{-1}x) = e(t)x \in \lambda(C)$. According to the assumption, $e(t)x = x$. So $x \in \lambda(C)$ which means that $\lambda(C)$ is closed, that is, λ is a closed map. □

Let (X, T, λ) be a T-space. If Y is a subset of X, then

$$TY := \{ty| \quad t \in T \quad \text{and} \quad y \in Y\}.$$

Y is called *invariant* under T if $TY = Y$.

Corollary 2.13. *Let (X, T, λ) be a T-space. Moreover T be compact, $Y \subseteq X$ and for each $x \in X$, $e(T) \subseteq T_x$, then*

(i) TY is closed if Y is closed;

(ii) TY is compact if Y is compact.

Theorem 2.14. *Let (X, T, λ) be a T-space and for each $x \in X$, $e(T) \subseteq T_x$. Then the projection map $\pi : X \to X/_T$ is an open map.*

Proof. To prove that λ is an open map, suppose $Y \subseteq X$ is an arbitrary open subset of X. We have

$$TY = \{ty \mid t \in T \text{ and } y \in Y\} = \pi^{-1}(\pi(Y)) = \bigcup_{t \in T} tY. \tag{4}$$

Since θ_t is a homeomorphism, then it is an open map. So for each $t \in T$, $\theta_t(Y) = tY$ is an open set in X. Hence $\pi^{-1}(\pi(Y))$ is an open set in X (from (4)). So $\pi(Y)$ is open in $X/_T$, which means that π is an open map. □

Definition 2.15. Suppose (X, T, λ) is a T-space. A generalized action λ is called *perfect* if $e(T) \subseteq T_x$ for each $x \in X$. In this sense, the T-space is called *perfect*.

Suppose that X and Y are two topological spaces. A mapping $f : X \to Y$ is called proper if for each compact subset A of Y, $f^{-1}(A)$ is a compact subset of X [15]. We know that if the mapping $f : X \to Y$ is closed and $f^{-1}(y)$ is compact for each $y \in Y$, then f is a proper map [15].

Theorem 2.16. *Let (X, T, λ) be a T-space, T be compact and λ be perfect. Then*

(i) *The projection map $\pi : X \to X/_T$ is a closed map;*

(ii) *The quotient space $X/_T$ is Hausdorff;*

(iii) *The projection map $\pi : X \to X/_T$ is a proper map;*

(iv) *X is compact if and only if $X/_T$ is compact;*

(v) *X is locally compact if and only if $X/_T$ is locally compact.*

Proof. (i) Suppose that $Z \subseteq X$ is an arbitrary closed subset of X. As λ is perfect and T is compact, according to Theorem 2.13, $TZ = \pi^{-1}(\pi(Z))$ is closed in X. Since π is surjective map, we can see that

$$\pi^{-1}(X/_T - \pi(Z)) = X - \pi^{-1}(\pi(Z)). \tag{5}$$

Since $\pi^{-1}(\pi(Z))$ is closed in X, then $X - \pi^{-1}(\pi(Z))$ is open in X. $X/_T - \pi(Z)$ is open in $X/_T$. Thus $\pi(Z)$ is closed in $X/_T$, hence π is a closed map.

(ii) Suppose that $[x]$ and $[y]$ are two distinct elements in $X/_T$. Clearly, we can see that $T(x) \cap T(y) = \varnothing$. As mentioned, $T(x) = \rho_x(T)$, that is, the T-orbit $T(x)$ is the image of T. But we know that ρ_x is continuous and T is compact, which imply that $T(x) = \pi^{-1}([x])$ is compact in X. In the same way, $T(y)$ is compact. Since X is Hausdorff and $T(x)$ is compact in X and $y \notin T(x)$, thus there exist disjoint open subsets U and V of X containing $T(x)$ and y, respectively [15]. Especially, it is easy to see that $T(x) \cap \overline{V} = \varnothing$. So $[x] = \pi(x) \notin \pi(\overline{V})$. On the other hand, $\pi(V)$ is open in $X/_T$. $\pi(V)$ contains of $[y]$. Moreover; since $\overline{V}$ is closed, $\pi(\overline{V})$ is closed in $X/_T$. Thus $X/_T - \pi(\overline{V})$ is an open subset that contains $[x]$. Therefore $X/_T - \pi(\overline{V})$ and $\pi(V)$ are disjoint open sets of $X/_T$ containing $[x]$ and $[y]$, respectively.

(iii) In (i), we saw that π is closed map and also in (ii), we shown that $T(x) = \pi^{-1}([x])$ is compact for each $x \in x$. So π is a proper map.

(iv) This follows easily from continuity of π and (iii).

(v) This follows from Theorem 2.14 and (iii).

□

Definition 2.17. Let (X, T, λ) be a T-space. The generalized action λ is called *proper* if the map $\hat{\lambda} : T \times X \to X \times X$ defined by $\hat{\lambda}(t, x) = (tx, x)$ is a proper map.

Theorem 2.18. *Let (X, T, λ) is a T-space. The generalized action λ is proper if and only if for every compact subset Y of X, the set $Y_T = \{t \in T \mid tY \cap Y \neq \varnothing\}$ is compact.*

Proof. Suppose that the mapping $\hat{\lambda}$ is proper and $Y \subset X$ is compact. We see that $Y_T = P(\hat{\lambda}^{-1}(Y \times Y))$ where $P : T \times X \to T$ is the projection map. So Y_T is compact. Conversely, Suppose that Y_T is compact, where Y is a compact subset of X. If $Z \subset X \times X$ is compact, and $Y := \pi_1(Z) \cup \pi_2(Z) \subset X$, where $\pi_1, \pi_2 : X \times X \to X$ are the projections on the first and second components, respectively, then Y is compact. Moreover $\hat{\lambda}^{-1}(Z) \subset \hat{\lambda}^{-1}(Y \times Y) \subset Y_T \times Y$. Since $X \times X$ is Hausdorff and Z is compact, then Z is a closed subset of $X \times X$. So $\hat{\lambda}^{-1}(Z)$ is closed by continuity and it is also a closed subset of the compact set $Y_T \times Y$. Thus Z is compact and λ is proper. □

Corollary 2.19. *Let (X, T, λ) be a T-space. If T is compact, then the generalized action λ is proper.*

Corollary 2.20. *Let (X, T, λ) be a T-space. If λ is proper, then*

(i). The stabilizer T_x is compact, where $x \in X$;

(ii). The orbit map ρ_x is proper, where $x \in X$.

Corollary 2.21. *Let (X, T, λ) be a T-space. If λ is proper, then $T(x)$ is closed subset of X and the quotient space $X/_T$ is Hausdorff.*

Example 2.22. Let $X = \mathbb{R}^2$ and let T be the generalized group of Example 2.3 which acts on X by

$$(r_1 e^{i\theta_1}).(r_2 e^{i\theta_2}) = r_1 r_2 e^{i\theta_2}.$$

The equivalence class $[x]$ is the set all $re^{i\theta_0}$ such that $r > 0$, where $x = r_0 e^{i\theta_0} \in \mathbb{R}^2 - \{(0, 0)\}$ and $[(0, 0)] = \{(0, 0)\}$. Moreover, for $x = r_1 e^{i\theta_1}$ and $y = r_2 e^{i\theta_2}$, $[x] = [y]$ if and only if $\theta_1 = \theta_2$. So $Y = X/_T \simeq S^1 \bigcup \{(0, 0)\}$ which is not a Hausdorff space with the quotient topology.

3 Maps On T-spaces

Now, let us consider the maps of T-spaces. First, we recall the notion of transitivity of maps on a topological space. Let X be a topological space and let $f : X \to X$ be a continuous mapping. Then the mapping f is called *(topologically) transitive* [11] if for every pair of non-empty open subsets U and V of X, there is some $n \in \mathbb{N}$ such that $f^n(U) \cap V \neq \varnothing$. Moreover, if X is compact, then f is transitive if and only if f is onto and there is a point in X with dense orbit [7].

Definition 3.1. Let (X, T, λ) be a T-space. Suppose $f : X \to X$ is a continuous map. The map f is called *T-transitive* if for every pair of non-empty open subsets U and V in X, there is some positive integer n and $t \in T$ such that $tf^n(U) \cap V \neq \varnothing$.

We can see that if λ is not trivial and f is a transitive map, then f is T-transitive. But the fallowing example shows that each T-transitive map need not be transitive.

Example 3.2. Let X be the topological space $[-1, 1]$ with the topology generated by Euclidean Metric. The topological generalized group $T = \{\pm 1\}$ with the multiplication $s \cdot t = s|t|$ acts on X by the generalized action $tx = t|x|$, where $t \in T$ and $x \in X$. Then $f : X \to X$ is defined by

$$f(x) = \begin{cases} \sqrt{x} & \text{if } \ 0 \leq x \leq 1 \\ -\sqrt{|x|} & \text{if } \ -1 \leq x \leq 0 \end{cases}$$

We can see that f is T-transitive but it is not transitive.

Definition 3.3. Let (X, T, λ) and (Y, T, μ) be two T-spaces. A continuous map $f : X \to Y$ is called

(i) *T-equivariant* if $f(\lambda(t, x)) = \mu(t, f(x))$ for $t \in T$ and $x \in X$, briefly, $f(tx) = tf(x)$;

(ii) *T-pseudoequivariant* if $f(T(x)) = T(f(x))$ for $x \in X$.

If $f : X \to Y$ is a T-equivariant between two T-spaces X and Y, then $T_x \subseteq T_{f(x)}$ for each $x \in X$. Moreover, an T-equivariant map is clearly T-pseudoequivariant but the converse is not true. The following example shows this.

Example 3.4. Let X and Y be two Euclidean space $\mathbb{R}$ and $T = \{\pm 1\}$. Then T with the multiplication $s \cdot t = s|t|$ is a topological generalized group. We define actions $\lambda, \theta : T \times \mathbb{R} \to \mathbb{R}$ by $\lambda(t, x) = t|x|$ and $\theta(t, x) = tx$, respectively. Now, we see that the identity map on $\mathbb{R}$ is a T-pseudoequivariant map which is not T-equivariant.

Theorem 3.5. *Let (X, d) be a compact metric space with no isolated point and let (X, T, λ) be a T-space which λ is perfect. Suppose $f : X \to X$ be a T-pseudoequivariant onto map. Then f is T-transitive if and only if there exists $x \in X$ such that $\{t \cdot f^n(x) \mid t \in T, n > o\}$ is dense in X.*

Proof. Suppose f is T-transitive. By using of Baire category theorem and the fact that λ is perfect, we can see that there is some x such that $\{t \cdot f^n(x) \mid t \in T, n > o\}$ is dense in X. Conversely, suppose that $\{tf^n(x) \mid t \in T, n > o\}$ is dense in X for some $x \in X$. Let U and V be two non-empty open subsets of X. Then there are $s, t \in T$ and $m, n \in \mathbb{N}$ such that $sf^m(x) \in U$ and $tf^n(x) \in V$. We assume that $m < n$. Since λ is perfect then $x \in f^{-m}(s^{-1}U)$. Thus $f^n(x) \in f^{n-m}(s^{-1}U)$. Moreover, since f is T-pseudoequivariant, then there is $r \in T$ such that $f^{n-m}(s^{-1}U) = rf^{n-m}(U)$. So $f^n(x) \in rf^{n-m}(U)$ and $tf^n(x) \in trf^{n-m}(U)$. Therefore, $tf^n(x) \in t^{'}f^p(U) \cap V$ where $t^{'} = tr \in T$ and $p = n - m \in \mathbb{N}$. So f is T-transitive. □

It may be that T-transitivity is not preserved by topological conjugacy. Now we introduce topological T-conjugacy that can preserve T-transitivity.

Definition 3.6. Let (X, T, λ) and (Y, T, θ) be two T-spaces. Moreover let $f : X \to X$ and $g : Y \to Y$ be two continuous maps. We say f is *topologically T-conjugate* to g if there is a T-pseudoequivariant homeomorphism $h : X \to Y$ such that $h \circ f = g \circ h$.

Theorem 3.7. *Let (X, T, λ) and (Y, T, θ) be two perfect T-spaces. If the continuous mapping $f : X \to X$ is topologically T-conjugate to the continuous mapping $g : Y \to Y$ then f is T-transitive if and only if g is T-transitive.*

Now, we recall that a *top sapce* T is a Hausdorff d-dimensional differentiable manifold which is also a topological generalized group such that mappings m_1 and m_2 are smooth [11].

Theorem 3.8. *Let X and Y be smooth manifolds and let T be a top space. Suppose $F : X \to Y$ is a smooth map that is equivariant respect to a transitive perfect smooth generalized action λ of T on X and a perfect smooth generalized action θ of T on Y. Then F has constant rank.*

Proof. Let $x \in X$. Since λ is transitive then for any $y \in X$, there is some $t \in T$ such that $tx = y$. Since $\theta_t \circ F = F \circ \lambda_t$ for each $t \in T$, then we have $\theta_{t*} \circ F_* = F_* \circ \lambda_{t*}$. The generalized actions λ and θ are perfect, then λ_{t*} and θ_{t*} are isomorphism. So the rank F at y is the same as its rank at x. Thus F has constant rank. □

References

[1] Ahmadi S.A. *Generalized Topological Groups and Genetic Recombination*, Journal of Dynamical Systems and Geometric Theories 2013; 11, 1: 51-58.

[2] Araujo J, Konieczny J. *Molaei's Generalized Groups are Completely Simple Semigroups*, Buletinul Institului Polithnic Din Iasi 2004; 48, 52: 1-5.

[3] Donmez D. *On the Validity of the Borel-Hirzebruch Formula for Topological Actions*, Turkish Journal of Mathematics 1998; 22: 193-198.

[4] Ebrahimi N. *Left Invariant Vector Fields of a Class of Top Spaces*, Balkan J. Geom. Appl. 2012; 14: 37-44.

[5] Farhangdoost M.R. *Action of Generalized Groups on Manifolds*, Acta Math. Univ. Comenianae. 2011; LXXX, 2: 229-235.

[6] Guner N. *A Borsuk-Ulak Theorem for Heisenberg Group Actions*, Turkish Journal of Mathematics 2000; 24: 345-357.

[7] Kolyada S, Snoha L. *Some Aspects of Topological Transitivity A Survey*, Iteration theory, ECIT 94, Opava, Grazer Math. Ber. 334, Karl Franzens Univ. Graz 1997; 3-35.

[8] Molaei M.R. *Generalized Actions*, Proceedings of the First International Conference on Geometry, Integrability and Quantization, Coral Press Scientific Publishing, September 1999; 1-10: 175-180.

[9] Molaei M.R. *Generalized Groups*, Buletinul Institului Polithnic Din Iasi 1999; Tomul XLV(XLIX): 21-24.

[10] Molaei M.R. *Topological Generalized Groups*, International Journal of Applied Mathematics 2000; 2, 9: 1055-1060.

[11] Molaei M.R. *Top Spaces*, J. Interdiscip. Math. 2004; 7, 2: 173-181.

[12] Molaei M.R. *Mathematical Structures Based on Completely Simple Semigroups*, Hadronic press, 2005.

[13] Molaei M.R. *Complete Semi-dynamical Systems*, Journal of Dynamical Systems and Geometric Theories 2005; 3, 2: 95-107.

[14] Molaei M.R, Khadekar G.S, Farhangdoost M.R. *On Top Spaces*, Balkan J. Geom. Appl. 2009; 11, 1: 101-106.

[15] Munkres J.R. Topology A First Course, Prentice Hall of India, 1975.